

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/293191297>

Low chronic workload and the acute: Chronic workload ratio are more predictive of injury than between-match recovery time: A two-season prospective cohort study in elite rugby leag...

Article in *British Journal of Sports Medicine* · February 2016

DOI: 10.1136/bjsports-2015-095364

CITATION

1

READS

731

5 authors, including:



Billy Hulin

University of Queensland

8 PUBLICATIONS 49 CITATIONS

SEE PROFILE



Tim Gabbett

235 PUBLICATIONS 5,786 CITATIONS

SEE PROFILE



John A Sampson

University of Wollongong

27 PUBLICATIONS 58 CITATIONS

SEE PROFILE

Low chronic workload and the acute:chronic workload ratio are more predictive of injury than between-match recovery time: a two-season prospective cohort study in elite rugby league players

Billy T Hulin,^{1,2} Tim J Gabbett,^{3,4} Peter Caputi,⁵ Daniel W Lawson,⁶ John A Sampson¹

¹Centre for Human and Applied Physiology, School of Medicine, University of Wollongong, Wollongong, New South Wales, Australia

²Football Department, St. George Illawarra Dragons Rugby League Football Club, Wollongong, New South Wales, Australia

³School of Exercise Science, Australian Catholic University, Brisbane, Queensland, Australia

⁴School of Human Movement Studies, University of Queensland, Brisbane, Queensland, Australia

⁵School of Psychology, University of Wollongong, Wollongong, New South Wales, Australia

⁶Baimed Physiotherapy and Sports Injury Clinic, Wollongong, New South Wales, Australia

Correspondence to

Billy T Hulin, Football Department, St. George Illawarra Dragons RLFC, 1/5 Burelli Street, Wollongong, NSW 2500, Australia; billyhulin@hotmail.com

Accepted 18 January 2016

ABSTRACT

Background Between-match recovery time, and acute and chronic workloads likely affect subsequent match-injury risk in elite rugby league players.

Methods Workloads of 28 players throughout two seasons were calculated during short (<7 days), and long (≥ 7 days) between-match recovery times. 'Acute' workloads (1 week) greater than 'chronic' workloads (4-week rolling average acute workload) resulted in acute:chronic workload ratios above 1.

Results No difference was found between the match-injury risk of short and long between-match recovery periods ($7.5 \pm 2.5\%$ vs $6.8 \pm 2.5\%$). When players had a short recovery between matches, high chronic workloads (18.9–22.0 km) were associated with a smaller risk of match injury than chronic workloads <18.9 km (relative risk (RR) range 0.27–0.32 (CI 0.08 to 0.92); likelihood range 90–95%, likely). Players who had shorter recovery and acute:chronic workload ratios ≥ 1.6 , were 3.4–5.8 times likely to sustain a match injury than players with lower acute:chronic workload ratios (RR range 3.41–5.80 (CI 1.17 to 19.2); likelihood range 96–99%, very likely). Acute:chronic workload ratios between 1.2 and 1.6 during short between-match recovery times demonstrated a greater risk of match injury than ratios between 1.0 and 1.2 (RR=2.88 (CI 0.97 to 8.55); likelihood=92%, likely).

Conclusions Contrary to the philosophy that high workloads and shorter recovery equate to increased injury risk, our data suggest that high and very-high chronic workloads may protect against match injury following shorter between-match recovery periods. Acute:chronic workload ratios ~ 1.5 are associated with a greater risk of match injury than lower acute:chronic workload ratios. Importantly, workloads can be manipulated to decrease the match-injury risk associated with shorter recovery time between matches.

INTRODUCTION

The likelihood of sustaining an injury in team sport can be influenced in part by: (1) the workloads that athletes are subjected to,^{1–3} or (2) the recovery time that is provided between matches.^{4–5} Three to 5 days recovery is required for match-induced reductions in neuromuscular and endocrine function to return to prematch values in team sport athletes.^{6–7} During the shortest between-match recovery times in rugby league (5 and 6 days), training and preparation between matches is congested and there is a greater incidence of injury

than during longer between-match recovery times (7 or more days).⁵ However, at least two studies failed to find any association between congested fixture periods and higher injury incidence in other team sports.^{8–9}

The equivocal evidence in relation to shorter between-match recovery times and increased injury risk may be due to the fact that previous studies have not investigated player workloads between matches.^{4–5–8–9} Higher workloads can occur during longer between-match recovery times¹⁰ and higher workloads^{1–3} or abrupt increases in workloads^{2–3–11} can increase injury risk. However, no study has investigated the combined influence of workloads and between-match recovery time on the risk of subsequent match injury in elite team sport athletes.

Regardless of the recovery time between matches, higher acute (ie, 1 week) workloads are associated with winning more matches in an elite team sport.¹² However, when acute workload exceeded chronic workload (ie, 4-week average acute workload), resulting in an acute:chronic workload ratio greater than 1, the probability of losing matches increased.¹² Furthermore, Hulin *et al*¹¹ recently demonstrated that acute and chronic workloads can have both positive and negative influences on injury risk in elite rugby league players. When players had an acute:chronic workload ratio close to 1, higher chronic workloads were associated with a smaller risk of injury than lower chronic workloads. Conversely, a large spike in acute workload relative to chronic workload (ie, ratios of >1.5) resulted in a threefold to fivefold increase in injury risk.^{2–11}

The effect of: (1) higher chronic workloads or (2) abrupt increases in acute workload, on the risk of sustaining a match injury following different between-match recovery times is currently unknown. Therefore, we modelled acute and chronic workloads with the risk of sustaining a match injury following short (5 and 6 days) and long (7, 8 and 9 days) between-match recovery times in elite rugby league players.

METHODS

Participants

Twenty-eight players (mean \pm SD; age, 24.8 \pm 3.4 years; height, 184.5 \pm 4.9 cm; body mass, 99.5 \pm 8.0 kg) from one elite rugby league club

To cite: Hulin BT, Gabbett TJ, Caputi P, *et al*. *Br J Sports Med* Published Online First: [please include Day Month Year] doi:10.1136/bjsports-2015-095364

participated in this study over two Australian National Rugby League (NRL) seasons. This study was conducted over the competition phase of each season (2×27 weeks). Only players participating in first-team matches and completing workloads between those matches were included in the analysis. Players received a clear explanation of the study and written consent was obtained. Experimental procedures were approved by the Institutional Review Board for Human Investigation.

Injury definition

Injury records were updated and maintained by the club's senior physiotherapist. An injury was defined as any time-loss injury that resulted in a player being unable to complete full training, or missing match time.^{11 13 14} Only injuries that were sustained in a match or reported subsequent to a match, prior to the next training session, were included in the combined analysis of workload and between-match recovery time.

Workload and between-match recovery time analysis

Workload was defined as absolute total distance (m) covered during all field training sessions and matches and was measured via GPS (GPSports, SPI-HPU 5 Hz (interpolated 15 Hz), Canberra, Australia). The GPS equipment used in this study has demonstrated adequate accuracy and reliability for measuring total distance covered^{15 16}; however, this equipment is not accurate or reliable when measuring changes in velocity,^{15 17} high-speed running^{15 18} and collisions.¹⁹ Therefore, these variables were excluded from analysis. In the event that a player did not wear a GPS unit or the GPS unit failed to collect data (<1% of the dataset), the player was given either the average workload of that training session or their average match workload over the season.²⁰

Between-match workloads were calculated as the total distance covered during all field training sessions and the following match. Data were categorised into between-match recovery times comparable with previous studies.^{5 10} Specifically, these recovery times consisted of <7 days (5 and 6 days, also referred to as 'short') between matches, or ≥7 days (7, 8 and 9 days, also referred to as 'long') between matches. Chronic workloads and acute:chronic workload ratios were calculated in accordance with previous work.^{2 11 21 22} Briefly, data from Monday to Sunday represented the acute workload, while the 4-week rolling average acute workload represented the chronic workload. An acute workload greater than chronic workload was indicated by an acute:chronic workload ratio greater than 1, and vice versa.^{2 11 21 22}

Workload data for short, and long between-match recovery times were further categorised into very-low through very-high categories according to percentile rank. These categories are displayed in [table 1](#). Injury-workload relationships were calculated among very-low through very-high between-match workloads, chronic workloads and acute:chronic workload ratios.

Statistical analysis

Match-injury risks were calculated as the total number of match injuries sustained relative to the total number of exposures to each workload classification.^{2 11 23} Null hypothesis testing was conducted using a binary logistic regression model with injury/no injury as the dependent variable. Between-match workloads, chronic workloads and acute:chronic workload ratios were independently modelled as predictor variables. Relative risk (RR) and 90% CI (lower-upper) were calculated to determine which workload variables increased or decreased the risk of match injury.²³ A RR of greater or less than 1 implied an increased or decreased risk of injury, respectively.

Table 1 Description of between-match workloads, chronic workloads and acute:chronic workload ratios during short and long between-match recovery times

| | Percentile rank (%) | Between-match recovery time | |
|-------------------------------------|---------------------|-----------------------------|-----------|
| | | Short | Long |
| Between-match workload (km) | | | |
| Very-low | ≤5 | ≤9.0 | ≤12.0 |
| Low | ≤25 | 9.1–12.5 | 12.1–16.2 |
| Moderate-low | <50 | 12.6–15.5 | 16.3–18.8 |
| Moderate-high | ≥50 | 15.6–17.8 | 18.9–21.6 |
| High | ≥75 | 17.9–21.4 | 21.7–25.3 |
| Very-high | ≥95 | ≥21.5 | ≥25.4 |
| Chronic workload (km) | | | |
| Very-low | ≤5 | ≤10.5 | ≤11.4 |
| Low | ≤25 | 10.6–14.3 | 11.5–14.7 |
| Moderate-low | <50 | 14.4–16.5 | 14.8–17.0 |
| Moderate-high | ≥50 | 16.6–18.8 | 17.1–20.0 |
| High | ≥75 | 18.9–22.0 | 20.1–23.0 |
| Very-high | ≥95 | ≥22.1 | ≥23.1 |
| Acute:chronic workload ratio | | | |
| Very-low | ≤5 | ≤0.66 | ≤0.80 |
| Low | ≤25 | 0.67–0.86 | 0.81–0.99 |
| Moderate-low | <50 | 0.87–1.01 | 1.00–1.09 |
| Moderate-high | ≥50 | 1.02–1.22 | 1.10–1.20 |
| High | ≥75 | 1.23–1.61 | 1.21–1.49 |
| Very-high | ≥95 | ≥1.62 | ≥1.50 |

The influence of between-player heterogeneity on the number of injuries sustained was examined using a Poisson regression model. In this model, positional group (forwards=1, backs=2) was used as a factor, while age, body mass and height were used as covariates.

Statistical analysis was performed in a similar fashion to other investigations of acute and chronic workloads and injury risk.¹¹ The p value derived from binary logistic regression and the value of the RR between groups were entered into a customised spreadsheet, which calculates 90% CI and the probabilities that the true effect was harmful, trivial and beneficial.^{24 25} These values were reported in quantitative and qualitative terms according to the following: ≥1%, very unlikely; ≥5%, unlikely; ≥25%, possibly; ≥75%, likely; ≥95%, very likely.^{24–26} Practical significance occurred when the probability that the true effect was either harmful or beneficial was ≥75%, likely.²⁴

RESULTS

There were 44 match, and 9 training injuries. No difference was found between the risk of match injury following short (injury risk=7.7±2.5%) and long (injury risk=6.8±2.5%) between-match recovery periods ([figure 1](#)). However, 8 of the 9 training injuries occurred when recovery time was <7 days (short).

The Poisson regression model did not demonstrate any relationship among the number of injuries sustained and positional group (p=0.303), age (p=0.325), height (p=0.221) or body mass (p=0.167).

Between-match workloads

[Figure 2A](#) depicts the risk of match injury with very-low through very-high between-match workloads during short, and long between-match recovery times. During short between-match recovery times, a high between-match workload (17.9–21.5 km) was associated with a risk of match-injury that was:




Figure 1 Risk of match injury \pm 90% CI subsequent to short (5 and 6 days) or long (7, 8 and 9 days) recovery time between matches.

(1) 87% smaller than low between-match workloads (RR=0.13 (CI 0.02 to 0.67); likelihood=97%, very likely), (2) 82% lower than moderate-low between-match workloads (RR=0.18 (CI 0.03 to 0.95); likelihood=94%, likely), and (3) 85% lower than very-high between-match workloads (RR=0.15 (CI 0.02 to 1.03); likelihood=94%, likely).

Chronic workloads

No match injuries were sustained when players (n=17) had a very-high chronic workload (≥ 22.1 km) during short between-match recovery times. A high chronic workload (18.9–22.1 km) was associated with a risk of match injury that was: (1) 73% lower than a moderate-low chronic workload (RR=0.27 (CI 0.08 to 0.92); likelihood=95%, very likely), and (2) 68% lower than a low chronic workload (RR=0.32 (CI 0.08 to 1.22); likelihood=90%, likely; figure 2B). During short between-match recovery times, there was a linear trend for *lower* risk of subsequent match injury as chronic workload *increased* (figure 2B).

Acute:chronic workload ratios

A high acute:chronic workload ratio (1.23–1.61) during short between-match recovery times demonstrated a risk of match injury that was 2.88 times greater than a moderate-high acute:chronic workload ratio combined with short recovery between matches (RR=2.88 (CI 0.97 to 8.55); likelihood=92%, likely; figure 2C). The risk of match injury with a very-high acute:chronic workload ratio (≥ 1.62) combined with short recovery between matches was: (1) 5.80 times greater than a moderate-high acute:chronic workload ratio (RR=5.80 (CI 1.75 to 19.2); likelihood=99%, very likely), and (2) 3.41 times greater than a low acute:chronic workload ratio (RR=3.41 (CI 1.17 to 9.91); likelihood=96%, very likely; figure 2C).

Long between match recovery times, combined with a very-high acute:chronic workload ratio (≥ 1.50) displayed a risk of match injury that was 4.46 times greater than a moderate-high acute:chronic workload ratio (RR=4.46 (CI 0.91 to 21.91); likelihood=92%, likely; figure 2C).

With the exception of high acute:chronic workload ratios, when players' acute:chronic workload ratios were taken into account, the recovery time between matches did not have a significant independent effect on injury risk in the subsequent match (figure 2C).

DISCUSSION

This study investigated the combined influence of recovery time between matches and player workloads on subsequent match-




Figure 2 Risk of match injury \pm 90% CI subsequent to short (5 and 6 days) or long (7, 8 and 9 days) recovery time between matches combined with between-match workloads (A), chronic workloads (B) and acute:chronic workload ratios (C). (a) Likely ($\geq 75\%$) different from moderate-high between-match workload and very likely ($\geq 95\%$) different from high between-match workload, during short between-match recovery times. (b) Likely ($\geq 75\%$) different from high between-match workload during short between-match recovery time. (c) Likely ($\geq 75\%$) different from high chronic workload during short between-match recovery time. (d) Likely ($\geq 75\%$) different from low, and moderate-high acute:chronic workload ratios during short between-match recovery time and high acute:chronic workload ratio during long between-match recovery time. (e) Very likely ($\geq 95\%$) different from low, and moderate-high acute:chronic workload ratio during either between-match recovery time, and high acute:chronic workload ratio during long between-match recovery time. (f) Likely ($\geq 75\%$) different from moderate-low acute:chronic workload ratio during either between-match recovery time and very-low and high acute:chronic workload ratio during short between match recovery time. (f) Likely ($\geq 75\%$) different from low, and moderate-high acute:chronic workload ratio during either between-match recovery time. Likely ($\geq 75\%$) different from high acute:chronic workload ratio during long between-match recovery time.

injury risk in elite rugby league. We found no difference in the risk of match injury, subsequent to short (7.7% injury risk) or long (6.8% injury risk) recovery between matches. Our findings

demonstrate that *higher* chronic workloads can *reduce* match injury risk following short between-match recovery times. However, increases in acute:chronic workload ratios are associated with a higher risk of match injury following short and long between-match recovery times.

The influence of workload and workload ratios on injury risk: does congestion matter?

We demonstrated that a controllable factor (workload) is associated with higher match-injury risk than non-modifiable between-match recovery time in isolation. Specifically, during short between-match recovery times, chronic workloads were associated with a linear trend for *lower* risk of injury as chronic workload *increased*. Additionally, high (1.2–1.6) and very-high (>1.6) acute:chronic workload ratios combined with a short recovery between matches were associated with a risk of match injury that was 2.8 and 5.8 times higher than an acute:chronic workload ratio between 1.0 and 1.2. Furthermore, with a long recovery between matches, very-high acute:chronic workload ratios (≥ 1.5) displayed a fourfold increase in match injury risk compared with ratios of 1.1–1.2. These findings demonstrate that *higher* workloads prior to short between-match recovery times can *decrease* injury risk, while sudden upgrades in workload will *increase* match-injury risk, regardless of the recovery time provided between matches.

In this study, the two greatest risks of match injury were observed when players had a very-high acute:chronic workload ratio combined with short (22% injury risk) or long (15% injury risk) recovery between matches. Additional match-injury risk factors in this study were when players had very-low to moderate-low chronic workloads during short between-match recovery times. Lower chronic workloads and sudden spikes in workload likely function together to increase the risk of match injury. That is, a high chronic workload may reduce the risk of injury due to the protection it provides against a very-high acute:chronic workload ratio. For example, chronic workloads of 19 and 12 km would be classed as high and low, respectively. Therefore, an acute workload greater than 20 km would result in moderate-high (ie, 1.1) and very-high (ie, 1.7) acute:chronic workload ratios for athletes with high and low chronic workloads, respectively. As such, higher chronic workloads may provide protection against a spike in acute workload, which was associated with the greatest risk of match injury in this study and overall injury in previous investigations.^{2 11}

Low through very-high between-match workloads demonstrated a slightly 'V'-shaped relationship during short between-match recovery times. Specifically, high between-match workloads were associated with a lower risk of match injury than low through moderate-high and very-high between-match workloads (figure 2A). These findings demonstrate that some, but not excessive, workload is essential to decrease the risk of match injury when short recovery has been provided between matches. Additionally, no injuries were sustained when players completed very-high workloads during long between-match recovery times. Collectively, these findings are in agreement with others—absolute acute workloads are not as predictive of injury as acute:chronic workload ratios.¹¹ Furthermore, previous research has shown that when acute:chronic workload ratios below 1 are prescribed, higher workloads between matches are associated with a greater number of victories in team sport.¹²

Practical implications for team coaches and strength/conditioning personnel

Collectively, these findings may be attractive to coaches and practitioners hoping to implement higher training workloads

between matches. However, attention to the myriad of factors other than workload, which are related to injury risk²⁷ and to the risk factors for overtraining and illness should also be considered when planning and prescribing between-match workloads.²⁸ For example, long-term excessive workload can cause non-functional over-reaching, characterised by decreases in performance and vigour and increased fatigue.^{28 29}

Previous studies may support our findings that a high chronic workload can reduce the risk of match injury. Specifically, Gabbett *et al* demonstrated that rugby league players with greater aerobic capacity,³⁰ and prolonged high-intensity running ability³¹ have a lower risk of injury. Furthermore, only 2 weeks of low-volume sprint interval training is required to elicit improvements in the high-intensity, intermittent running ability and aerobic capacity of team sport athletes.³² Therefore, considering that rugby league training and match play involves sprinting efforts interspersed with low-intensity activity,^{33 34} the athletes in the current study that achieved a higher chronic workload may have improved or maintained these physical qualities and in turn decreased their risk of sustaining a match injury when provided a short recovery between matches. However, for chronic workloads to be increased and provide resistance to injury, an acute:chronic workload ratio greater than 1 must be cautiously prescribed at strategic periods throughout the season. The findings of the present study suggest that the associated injury risk would be lower if workloads were increased during between-match recovery times of 7 or more days, with an acute:chronic workload ratio between 1 and 1.5.

Limitations and future directions

Valid analysis of injury risk in relation to collisions, high-speed running and accelerations is not possible with the GPS equipment used in this study.^{15–19} Injury risk in team sport athletes can also be attributed to multiple factors that have not been included in this study.^{23 27} However, to achieve the statistical power required to investigate other factors such as previous injury, age and physical fitness, in conjunction with the modifiable and non-modifiable injury risk factors in the present study, would require a considerably larger data set comprising many teams over a number of seasons.²³ Of course that brings with it the confound of different training regimens (an inherent limitation of 'ecological' research).

Summary and conclusion

Our data demonstrated that when short and long between-match recovery times are viewed in isolation, no difference in injury risk occurred. This study suggests that low chronic workloads, and the acute:chronic workload ratio are more predictive of injury than merely the recovery time provided between matches. Specifically, a higher chronic workload provides protection against a spike in acute workload, which was associated with the greatest risk of match injury in this study. Furthermore, provided that very-high acute:chronic workload ratios (~1.5) are avoided, higher between-match workloads can be achieved without increasing injury risk in elite rugby league players. This study offers practitioners fresh insight of how a controllable factor (workload) can be modified to decrease match-injury risk during elite rugby league competition.

Contributors BTH, TJG and JAS are responsible for the concept and design of this project. Injury and workload data were collected by DL and BTH, respectively. BTH was responsible for data analysis. PC revised this manuscript and provided statistical expertise.

What are the findings?

- ▶ Without taking other factors into account, the match-injury risk following 'short' between-match recovery times (5 or 6 days) was 7.7%; this was no different to the match-injury risk of 6.8% following 'long' between-match recovery times (≥ 7 days).
- ▶ Higher workloads prior to short between-match recovery times may decrease injury risk, while sudden upgrades in workload will likely increase match-injury risk, regardless of the recovery time provided between matches.
- ▶ This study adds to the body of literature suggesting that the acute:chronic workload ratio should be considered when evaluating workload-related injury risk in various sports.

How might it impact on clinical practice in the future?

- ▶ Using these findings to plan training across a season, taking into account between-match recovery times, may decrease injury risk.
- ▶ The positive and negative influences of workload on injury risk may be useful for strength and conditioning coaches and clinicians involved in planning training programmes in elite rugby league.

Funding Technical or equipment support for this study was not provided by any outside companies, manufacturers or organisations. BTH was funded by a postgraduate research scholarship supported by the University of Wollongong and the St. George Illawarra Dragons Rugby League Football Club.

Competing interests None declared.

Patient consent Obtained.

Ethics approval UOW and ISLHD Health and Medical Human Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Gabbett TJ, Jenkins DG. Relationship between training load and injury in professional rugby league players. *J Sci Med Sport* 2011;14:204–9.
- 2 Hulin BT, Gabbett TJ, Blanch P, et al. Spikes in acute workload are associated with increased injury risk in elite cricket fast bowlers. *Br J Sports Med* 2014;48:708–12.
- 3 Rogalski B, Dawson B, Heasman J, et al. Training and game loads and injury risk in elite Australian footballers. *J Sci Med Sport* 2013;16:499–503.
- 4 Dupont G, Nedelec M, McCall A, et al. Effect of 2 soccer matches in a week on physical performance and injury rate. *Am J Sports Med* 2010;38:1752–8.
- 5 Murray NB, Gabbett TJ, Chamari K. Effect of different between-match recovery times on the activity profiles and injury rates of national rugby league players. *J Strength Cond Res* 2014;28:3476–83.
- 6 Cormack SJ, Newton RU, McGuigan MR. Neuromuscular and endocrine responses of elite players to an Australian rules football match. *Int J Sports Physiol Perform* 2008;3:359–74.
- 7 McLellan CP, Lovell DI, Gass GC. Biochemical and endocrine responses to impact and collision during elite Rugby League match play. *J Strength Cond Res* 2011;25:1553–62.
- 8 Carling C, Le Gall F, Dupont G. Are physical performance and injury risk in a professional soccer team in match-play affected over a prolonged period of fixture congestion? *Int J Sports Med* 2012;33:36–42.
- 9 Dellal A, Lago-Peñas C, Rey E. The effects of a congested fixture period on physical performance, technical activity and injury rate during matches in a professional soccer team. *Br J Sports Med* 2015;49:390–4.
- 10 Moreira A, Kempton T, Aoki MS, et al. The impact of 3 different-length between-matches microcycles on training loads in professional rugby league players. *Int J Sports Physiol Perform* 2015;10:767–73.
- 11 Hulin BT, Gabbett TJ, Lawson DW, et al. The acute:chronic workload ratio predicts injury: high chronic workload may decrease injury risk in elite rugby league players. *Br J Sports Med* 2016;50:231–6.
- 12 Aughey R, Elias GP, Esmaeili A, et al. Does the recent internal load and strain on players affect match outcome in elite Australian football? *J Sci Med Sport* 2016;19:182–6.
- 13 Gabbett TJ. Influence of training and match intensity on injuries in rugby league. *J Sports Sci* 2004;22:409–17.
- 14 King DA, Hume PA, Milburn PD, et al. Match and training injuries in rugby league: a review of published studies. *Sports Med* 2010;40:163–78.
- 15 Petersen C, Pyne P, Portus M, et al. Validity and reliability of GPS units to monitor cricket-specific movement patterns. *Int J Sports Physiol Perform* 2009;4:381–93.
- 16 Rampinini E, Alberti G, Fiorenza M, et al. Accuracy of GPS devices for measuring high-intensity running in field-based team sports. *Int J Sports Med* 2015;36:49–53.
- 17 Buchheit M, Al Haddad H, Simpson BM, et al. Monitoring accelerations with GPS in football: time to slow down? *Int J Sports Physiol Perform* 2014;9:442–5.
- 18 Johnston RJ, Watsford ML, Kelly SJ, et al. Validity and interunit reliability of 10 Hz and 15 Hz GPS units for assessing athlete movement demands. *J Strength Cond Res* 2014;28:1649–55.
- 19 Gabbett TJ. Quantifying the physical demands of collision sports: does microsensor technology measure what it claims to measure? *J Strength Cond Res* 2013;27:2319–22.
- 20 Colby MJ, Dawson B, Heasman J, et al. Accelerometer and GPS-derived running loads and injury risk in elite Australian footballers. *J Strength Cond Res* 2014;28:2244–52.
- 21 Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016. Published Online First: 12 Jan 2016 doi:10.1136/bjsports-2015-095788
- 22 Gabbett TJ, Hulin BT, Blanch P, et al. High training workloads alone do not cause sports injuries: how you get there is the real issue. *Br J Sports Med* Published Online First: 21 Jan 2016 doi:10.1136/bjsports-2015-095567
- 23 Bahr R, Holme I. Risk factors for sports injuries—a methodological approach. *Br J Sports Med* 2003;37:384–92.
- 24 Hopkins WG, Marshall SW, Batterham AM, et al. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc* 2009;41:3–13.
- 25 Hopkins WG. A spreadsheet for deriving a confidence interval, mechanistic inference and clinical inference from a P value. *Sportscience* 2007;11:16–20.
- 26 Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. *Int J Sports Physiol Perform* 2006;1:50–7.
- 27 Hägglund M, Waldén M, Ekstrand J. Risk factors for lower extremity muscle injury in professional soccer: the UEFA injury study. *Am J Sports Med* 2013;41:327–35.
- 28 Meeusen R, Duclos M, Gleeson M, et al. Prevention, diagnosis and treatment of the overtraining syndrome. *Eur J Sport Sci* 2006;6:1–14.
- 29 Foster C, Lehmann M. Monitoring training in athlete with reference to overtraining syndrome. *Med Sci Sports Exerc* 1998;30:1164–8.
- 30 Gabbett TJ, Domrow N. Risk factors for injury in sub-elite rugby league players. *Am J Sports Med* 2005;33:428–34.
- 31 Gabbett TJ, Ullah S, Finch CF. Identifying risk factors for contact injury in professional rugby league players—application of a frailty model for recurrent injury. *J Sci Med Sport* 2012;15:496–504.
- 32 Macpherson TW, Weston M. The effect of low-volume sprint interval training on the development and subsequent maintenance of aerobic fitness in soccer players. *Int J Sports Physiol Perform* 2015;10:332–8.
- 33 Gabbett TJ, Jenkins DG, Abernethy B. Physical demands of professional rugby league training and competition using microtechnology. *J Sci Med Sport* 2012;15:80–6.
- 34 Gabbett TJ. Sprinting patterns of National Rugby League competition. *J Strength Cond Res* 2012;26:121–30.



Low chronic workload and the acute:chronic workload ratio are more predictive of injury than between-match recovery time: a two-season prospective cohort study in elite rugby league players

Billy T Hulin, Tim J Gabbett, Peter Caputi, Daniel W Lawson and John A Sampson

Br J Sports Med published online February 5, 2016

Updated information and services can be found at:

<http://bjsm.bmj.com/content/early/2016/02/05/bjsports-2015-095364>

These include:

References

This article cites 32 articles, 7 of which you can access for free at: <http://bjsm.bmj.com/content/early/2016/02/05/bjsports-2015-095364#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Rugby](#) (161)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>